

# Lamivudine

**Brand Name:** Epivir



## Drug Description

Lamivudine, a synthetic antiretroviral agent, is a dideoxynucleoside reverse transcriptase inhibitor. Lamivudine is the negative enantiomer of a dideoxy analogue of cytidine and is structurally similar to zalcitabine (2',3'-dideoxycytidine, ddC). Lamivudine differs structurally from zalcitabine in that the 3'-carbon of the ribose ring is replaced with sulfur, forming an oxathiolane ring. The absence of a free 3'-hydroxy group on the oxathiolane ring results in the inability of lamivudine to form phosphodiester linkages at this position. Both the positive and negative enantiomers of 2',3'-dideoxy,3'-thiacytidine exhibit antiviral activity in vitro, but lamivudine appears to exhibit greater antiviral activity and to be considerably less cytotoxic than the positive enantiomer. [1]

## HIV/AIDS-Related Uses

Lamivudine was approved by the FDA on November 17, 1995, for use in combination with other antiretroviral agents for the treatment of HIV infection in adults and in pediatric patients 3 months of age and older.[2] [3] Lamivudine should always be used in conjunction with other antiretroviral agents and should not be used alone in the management of HIV infection. Lamivudine usually is used in three- or four-drug regimens that include another nucleoside reverse transcriptase inhibitor (NRTI) and either one or two protease inhibitors (PIs) or a nonnucleoside reverse transcriptase inhibitor (NNRTI).[4]

Lamivudine has been used in combination with zidovudine for prevention of maternal-fetal transmission of HIV. Although the safety and efficacy of this two-drug regimen has not been established, it is considered one of several options used in HIV-infected women in labor who have received no prior antiretroviral therapy. Lamivudine is also used in conjunction with zidovudine or, alternatively, with stavudine for postexposure prophylaxis of HIV infection in health care workers and other individuals exposed occupationally to blood, body fluids, or tissues associated with a risk for transmission of the HIV virus.[5]

## Non-HIV/AIDS-Related Uses

Lamivudine is used to treat chronic hepatitis B virus (HBV) infection associated with evidence of hepatitis B viral replication and active liver inflammation.[6] It is administered in doses lower than those used to treat HIV infection. The formulation and dosage of lamivudine used in HBV therapy are not appropriate for patients dually infected with HIV and HBV.[7] Patients with HIV infection should receive only dosing forms appropriate for treatment of HIV. The safety and efficacy of lamivudine have not been established for treatment of chronic HBV in patients dually infected with HIV and HBV.[8]

## Pharmacology

Lamivudine exerts a virustatic effect against retroviruses by acting as a reverse transcriptase inhibitor. Lamivudine is phosphorylated intracellularly to its active 5'-triphosphate metabolite, lamivudine triphosphate (L-TP, also known as 3TC-TP), which inhibits HIV reverse transcription and hepatitis B virus polymerase activity via viral DNA chain termination.[9] The principal mode of action of L-TP is inhibition of HIV reverse transcription via viral DNA chain termination after incorporation of the nucleoside analogue. L-TP is a weak inhibitor of mitochondrial DNA polymerase and mammalian DNA polymerases alpha and beta.[10] 3TC-LP is a structural analogue of deoxycytidine triphosphate (dC-TP), the natural substrate for reverse transcriptase. 3TC-TP appears to compete with naturally occurring dC-TP for incorporation into viral DNA by reverse transcriptase. Following incorporation of 3TC-TP into the viral DNA chain instead of dC-TP, viral DNA synthesis is terminated prematurely because the absence of a 3'-hydroxyl group on the oxathiolane ring prevents further 5' to 3' phosphodiester linkages. Lamivudine has in vitro virustatic activity against HIV-1, HIV-2, and human hepatitis B viruses, but it appears to be inactive against other common human viruses (e.g., cytomegalovirus, Epstein-Barr virus, influenza virus, herpes simplex virus types 1 and 2, respiratory syncytial virus, varicella-zoster virus).[11]

# Lamivudine

## Pharmacology (cont.)

Lamivudine is rapidly absorbed, with bioavailability from 80% to 88% in adults and adolescents and from 66% to 68% in children. Food delays the peak serum concentration; however, there is no significant difference in bioavailability when lamivudine is taken with food. Time to peak concentration is approximately 0.5 to 2 hours after a single 100-mg dose; with food, it increases to approximately 3.2 hours; with fasting, it is about 1 hour.[12]

Lamivudine is widely distributed after administration. Lamivudine crosses the blood-brain barrier and is distributed into the cerebrospinal fluid (CSF) to a limited extent. In children, CSF concentrations have ranged from 10% to 17% of the corresponding non-steady-state serum concentration.[13] Apparent volume of distribution after intravenous (IV) administration in 20 patients was 1.3 l/kg, suggesting that lamivudine distributes into extravascular spaces. The volume of distribution was independent of dose and did not correlate with body weight.[14]

Lamivudine is in FDA Pregnancy Category C. There are no adequate and well-controlled studies done in pregnant women; however, lamivudine has been found to cross the placenta in humans. Rats given doses of lamivudine up to 130 times the usual human dose based on body surface area revealed no evidence of impaired fertility. There was no evidence of impaired fertility or adverse effects on offspring in rats given doses up to 4000 mg per kg daily (plasma levels 80 to 120 times those in humans). Some evidence of embryo lethality in rabbits, but not in rats, was observed at doses similar to the usual human dose. To monitor maternal-fetal outcomes of pregnant women exposed to lamivudine and other antiretrovirals, the Antiretroviral Pregnancy Registry has been established. Physicians may register patients by calling 1-800-258-4263 or at <http://www.APRegistry.com>. Lamivudine has been found to distribute into breast milk in humans; nursing mothers are recommended by the manufacturer not to breast-feed while taking this medication.[15]

Plasma protein binding is low (36%).[16] [17]

Metabolism is a minor route of elimination. In humans, the only known metabolite is the trans-sulfoxide metabolite. Within 12 hours after a single oral dose in six HIV-infected adults, 5.2% was excreted as the trans-sulfoxide metabolite in the urine. Serum concentrations of this metabolite have not been determined. The majority of lamivudine is eliminated unchanged in urine by active organic cationic secretion. In 20 HIV-infected patients given a single IV dose, renal clearance was 280.4 +/- 75.2 ml/min, representing 71% +/- 16% of total clearance of the drug. In most single-dose studies in infected patients, the mean elimination half-life ranged from 5 to 7 hours. Oral clearance and elimination half-life were independent of dose and body weight over an oral dosing range of 0.25 mg/kg to 10 mg/kg.[18] The half-life of intracellular lamivudine triphosphate is 11 to 15 hours; serum half-life of lamivudine is about 2.6 hours in adults and 1.7 to 2 hours in children. The renal clearance of lamivudine is greater than the glomerular filtration rate, implying active secretion into the renal tubules. Hemodialysis increases lamivudine clearance by a mean of 24 ml/min, but the length of dialysis treatment (i.e., 24 hours) may not be long enough to alter mean lamivudine exposure. It is not known if lamivudine is removed by peritoneal dialysis or continuous (24 hour) hemodialysis.[19]

Resistance to lamivudine can be produced in vitro by serial passage of HIV-1 in the presence of increasing concentrations of the drug, and strains of HIV-1 with in vitro resistance to lamivudine have emerged during therapy with the drug. Primary infection with lamivudine-resistant HIV-1 has been reported rarely in adults who were treatment naive. While some strains of zidovudine-resistant HIV-1 may be susceptible to lamivudine, strains resistant to both zidovudine and lamivudine have been isolated. HIV isolates resistant to zalcitabine, zidovudine, didanosine, lamivudine, and stavudine have been isolated from a limited number of patients who received zidovudine in conjunction with zalcitabine or didanosine for 1 year or longer. Mutations identified in these multidrug-resistant isolates were Ala62 to Val, Val75 to Ile, Phe77 to Leu, Phe116 to Tyr, and Gln151 to Met; the mutation at position 151 appears to play an important role in the development of multidrug resistance. The possibility of cross resistance

# Lamivudine

## Pharmacology (cont.)

---

among lamivudine, didanosine, and zalcitabine based on reverse transcriptase codon 184 mutations also is of concern.[20]

## Adverse Events/Toxicity

---

Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues alone or in combination, including lamivudine and other antiretrovirals. Female gender, obesity, and prolonged exposure to antiretroviral nucleoside analogues may be risk factors. Such cases have occurred in patients with and without known risk factors for liver disease. Treatment with lamivudine should be suspended in any patient who develops clinical or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity (which may include hepatomegaly and steatosis even in the absence of marked transaminase elevations).[21] [22]

Post-treatment exacerbations of hepatitis B virus infection have been reported in non-HIV infected patients treated with lamivudine for chronic hepatitis B virus (HBV) infection when lamivudine therapy was discontinued. Similar exacerbations of HBV infection have been reported in patients infected with both HIV and HBV when lamivudine therapy was switched to a regimen not containing lamivudine. The causal relationship between discontinuation of lamivudine therapy and exacerbation of HBV infection is unknown. Patients should be closely monitored with both clinical and laboratory follow-up for at least several months after stopping treatment. There is insufficient evidence to determine whether reinitiation of lamivudine alters the course of post-treatment exacerbations of hepatitis.[23]

Adverse effects seen with the use of lamivudine include pancreatitis, paresthesia and peripheral neuropathy, or splenomegaly, and are more commonly observed in pediatric patients than in adults.[24]

## Drug and Food Interactions

---

Lamivudine and zalcitabine may inhibit the intracellular phosphorylation of one another. Consequently, lamivudine should not be coadministered with zalcitabine.[25]

Lamivudine exposure was increased by 44% and lamivudine renal clearance was decreased by 30% when coadministered with sulfamethoxazole/trimethoprim. Concurrent administration of lamivudine and zidovudine in one small study resulted in a 39% increase in peak plasma concentration of zidovudine with no significant changes in the area under the concentration-time curve (AUC) or total clearance of lamivudine or zidovudine.[26]

Concurrent administration of lamivudine with indinavir and zidovudine resulted in a 6% decrease in AUC of lamivudine, no change in AUC of indinavir, and a 36% increase in AUC of zidovudine. No adjustment in dose is necessary. Concurrent administration of lamivudine with drugs associated with pancreatitis (e.g., alcohol, didanosine, intravenous pentamidine, sulfonamides, and zalcitabine) or with drugs associated with peripheral neuropathy (dapsone, didanosine, isoniazid, stavudine, and zalcitabine) should be avoided or used with caution.[27]

The higher and lower dose formulations of lamivudine should not be used concurrently. Concurrent administration of products that also contain lamivudine, including the coformulations of lamivudine and zidovudine, and abacavir sulfate, lamivudine, and zidovudine, should be avoided.[28]

## Contraindications

---

Lamivudine is contraindicated in patients with previously demonstrated clinically significant hypersensitivity to any of the components of the products.[29]

Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues alone or in combination, including lamivudine and other antiretrovirals. Lamivudine for HIV in oral solution or tablet form contain higher doses of the active

# Lamivudine



## Contraindications (cont.)

---

ingredient (lamivudine) than lamivudine used to treat chronic hepatitis B. Patients with HIV infection should receive only dosing forms appropriate for treatment of HIV.[30]

In pediatric patients with a history of prior antiretroviral nucleoside exposure, a history of pancreatitis, or other significant risk factors for the development of pancreatitis, lamivudine should be used in caution. Treatment with lamivudine should be stopped immediately if clinical signs, symptoms, or laboratory abnormalities suggestive of pancreatitis occur.[31]

Risk-benefit should be considered in patients with HIV and/or renal function impairment.[32]

## Clinical Trials

---

For information on clinical trials that involve Lamivudine, visit the ClinicalTrials.gov web site at <http://www.clinicaltrials.gov>. In the Search box, enter: Lamivudine AND HIV Infections.

## Dosing Information

---

Mode of Delivery: Oral (oral solution and tablets).[33]

Dosage Form: For use primarily in hepatitis B virus treatment: 100 mg tablets and 5 mg/ml oral solution.[34]

For use primarily in HIV treatment: 150 mg tablets and 10 mg/ml oral solution.[35]

Fixed combination tablets with zidovudine (lamivudine 150 mg and zidovudine 300 mg per tablet).[36]

Fixed combination tablets with abacavir sulfate and zidovudine (lamivudine 150 mg, abacavir sulfate 300 mg, and zidovudine 300 mg per tablet).[37]

Storage: Lamivudine tablets should be stored at 25 C (77 F), with excursions permitted to 15 C to 30 C (59 F and 86 F). Lamivudine oral solution need not be reconstituted and should be stored at 25 C (77 F) in tightly closed bottles.[38]

## Chemistry

---

CAS Name: 2(1H)-Pyrimidinone, 4-amino-1-((2R,5S)-2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-[39]

CAS Number: 134678-17-4[40]

Molecular formula: C<sub>8</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>S[41]

C41.91%, H4.84%, N18.33%, O20.94%, S13.99%[42]

Molecular weight: 229.26[43]

Melting point: 160 to 162 C[44]

Physical Description: White to off-white crystalline solid.[45]

Stability: Lamivudine oral solution need not be reconstituted.[46]

Solubility: Approximately 70 mg/ml in water at 20 C.[47]

## Other Names

---

(-)-2'-Deoxy-3'-thiacytidine[48]

BCH-189[49]

GR-109714X[50]

3TC[51]

Epivir-HBV[52]

## Further Reading

---

Rockstroh JK. Management of hepatitis B and C in HIV co-infected patients. J Acquir Immune Defic Syndr. 2003 Sep;34 Suppl 1:S59-65. PMID: 14562859

Sullivan JL. Prevention of mother-to-child transmission of HIV--what next? J Acquir Immune Defic Syndr. 2003 Sep;34 Suppl 1:S67-72. PMID: 14562860

Thio CL. Management of chronic hepatitis B in the



# Lamivudine



## Further Reading (cont.)

---

HIV-infected patient. AIDS Read. 2004 Mar;14(3):122-9, 133, 136-7. Review. PMID: 15068000

## Manufacturer Information

---

Lamivudine  
GlaxoSmithKline  
5 Moore Drive  
Research Triangle Park, NC 27709  
(888) 825-5249

Epivir  
GlaxoSmithKline  
5 Moore Drive  
Research Triangle Park, NC 27709  
(888) 825-5249

## For More Information

---

Contact your doctor or an AIDSinfo Health Information Specialist:

- Via Phone: 1-800-448-0440 Monday - Friday, 12:00 p.m. (Noon) - 5:00 p.m. ET
- Via Live Help: [http://aidsinfo.nih.gov/live\\_help](http://aidsinfo.nih.gov/live_help) Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET

## References

---

1. AHFS Drug Information - 2003; p. 670
2. GlaxoSmithKline - Epivir Product Information, September 2003, p. 7. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
3. AHFS Drug Information - 2003; p. 665
4. AHFS Drug Information - 2003; p. 665
5. AHFS Drug Information - 2003; p. 666
6. GlaxoSmithKline - Epivir-HBV Product Information, September 2003, p. 7. Available at: [http://us.gsk.com/products/assets/us\\_epivir\\_hbv.pdf](http://us.gsk.com/products/assets/us_epivir_hbv.pdf). Accessed 07/01/04.
7. GlaxoSmithKline - Epivir-HBV Product Information, September 2003, p. 10. Available at: [http://us.gsk.com/products/assets/us\\_epivir\\_hbv.pdf](http://us.gsk.com/products/assets/us_epivir_hbv.pdf). Accessed 07/01/04.
8. GlaxoSmithKline - Epivir Product Information, September 2003, p. 12. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.

# Lamivudine



9. USP DI - 2003; p. 1646
10. GlaxoSmithKline - Epivir Product Information, September 2003, p. 2. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
11. AHFS Drug Information - 2003; p. 670
12. USP DI - 2003; p. 1647
13. USP DI - 2003; p. 1647
14. GlaxoSmithKline - Epivir Product Information, September 2003, pp. 5-6. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
15. USP DI - 2003; p. 1647
16. USP DI - 2003; p. 1647
17. USP DI - 2003; p. 1647
18. GlaxoSmithKline - Epivir Product Information, September 2003, pp. 5-6. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
19. USP DI - 2003; p. 1647
20. AHFS Drug Information - 2003; p. 671
21. GlaxoSmithKline - Epivir Product Information, September 2003, p. 11. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
22. USP DI - 2003; p. 1648
23. GlaxoSmithKline - Epivir Product Information, September 2003, p. 12. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
24. USP DI - 2003; p. 1648
25. GlaxoSmithKline - Epivir Product Information, September 2003, p. 13. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
26. USP DI - 2003; pp. 1647-1648
27. USP DI - 2003; pp. 1647-1648
28. USP DI - 2003; pp. 1647-1648
29. GlaxoSmithKline - Epivir Product Information, September 2003, p. 11. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
30. GlaxoSmithKline - Epivir Product Information, September 2003, p. 1. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
31. GlaxoSmithKline - Epivir Product Information, September 2003, p. 11. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
32. USP DI - 2003; p. 1648
33. AHFS Drug Information - 2003; p. 671
34. AHFS Drug Information - 2003; p. 671
35. AHFS Drug Information - 2003; p. 671
36. AHFS Drug Information - 2003; p. 671
37. AHFS Drug Information - 2003; p. 671
38. GlaxoSmithKline - Epivir Product Information, September 2003, p. 23. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
39. ChemIDplus - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 07/01/04.
40. ChemIDplus - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 07/01/04.

# Lamivudine



41. Merck Index - 2001; p. 958
42. Merck Index - 2001; p. 958
43. Merck Index - 2001; p. 958
44. Merck Index - 2001; p. 959
45. GlaxoSmithKline - Epivir Product Information, September 2003, p. 1. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
46. GlaxoSmithKline - Epivir Product Information, September 2003, p. 23. Available at: [http://us.gsk.com/products/assets/us\\_epivir.pdf](http://us.gsk.com/products/assets/us_epivir.pdf). Accessed 07/01/04.
47. Merck Index - 2001; p. 959
48. ChemIDplus - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 07/01/04.
49. MeSH - Available at: <http://www.nlm.nih.gov/mesh/Mbrowser.html>. Accessed 07/01/04.
50. ChemIDplus - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 07/01/04.
51. ChemIDplus - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 07/01/04.
52. ChemIDplus - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 07/01/04.